



Bark Grafting Pecans

Eric Stafne

Extension Horticulturist
Commercial Pecans and Tree Fruits

Becky Carroll

Senior Agriculturalist, Fruit Physiology

Since pecans will not reproduce true from seeds, growers can't just plant a good quality pecan nut and get a pecan tree like its parents. Instead, propagation of pecans requires grafting—an asexual propagation process that is really not difficult and often quite satisfying to those who try it. In fact, many persons who do this start with native or seedling pecans which grow by the thousands in creek and river valleys in 60 Oklahoma counties. By grafting a shoot (scion) containing buds from desirable stock to the native trees—they turn these less productive seedlings into some of the best adapted, productive trees.

The modified inlay bark graft process described here is one of the most convenient methods of pecan grafting. It basically involves cutting off most or the entire top of a small growing pecan tree, grafting desirable new graftwood at the cut, and then watching the tree produce a new top. This is done during April and May in Oklahoma, soon after growth starts and the bark begins to slip on the stock trees. The bark on the scion, or graftwood, must be tight, however. Stock for the grafts must be dormant when cut during the winter usually in late December through February and kept in cold storage until used.

Successful bud growth is on the way when callus tissue forms between the stock tree and the scion.

Definitions:

Topworking - A method of changing the top of a tree (usually fairly large) to a desired variety. This is commonly done by inlay bark grafting.

Scionwood - (Graftwood, Propagation wood) mature dormant shoots usually of the previous season's growth. Consult HLA-6217 "Collecting and Storing Pecan Propagation Wood."

Scion - A section of scionwood to be grafted onto a stock. It is from a known variety one or two years old, 3/8 to 1/2 inches in diameter and 5 to 6 inches long.

Stock - A root, trunk or limb on which the scion is grafted.

Cambium - A thin layer of cells located between the bark and wood; capable of dividing and forming new cells. For a successful graft union, it is essential that the cambium of the scion be placed in close contact with the cambium of the stock.

Callus - A mass of cells that develop from and around wounded plant tissue. It occurs at the junction of the graft union, arising from the cambium of both scion and stock.

Oklahoma Cooperative Extension Fact Sheets
are also available on our website at:
<http://osufacts.okstate.edu>

Grafting - The process of inserting the scion into the stock.

What you need:

Stock trees — Select healthy, vigorous trees measuring one to four inches in diameter where the scion, or graft, will be set.

Graftwood — cut your own scions from desirable varieties from dormant wood and keep in cold storage until used; or, see your county Cooperative Extension educator for sources where suitable graftwood can be purchased or online at <http://www.hortla.okstate.edu/pecan/images/graftwood.pdf>.


Tools — grafting knife; small, sturdy, coarse-toothed saw; hand shears for cutting scions and removing smaller branches; sturdy, large-handled knife; small hammer for driving nails without damaging bark; no. 18 nails, 5/8" to 3/4" long; carpenter's apron.

Supplies — household aluminum foil; quart-sized polyethylene bags; rubber bands; masking or grafting tape


Grafting Procedure

First prepare the stock by sawing off the trunk or limb straight across with a sharp saw. Make this cut immediately above a 6" to 10" straight smooth section of the stock at a convenient working height, (usually 4 or 5 feet). Where cattle or horses are in the area, make the cut 6 to 8 feet above the ground. If available, leave one to three branches below the cut to keep the tree actively growing and for future use as a "throttle" in regulating the growth of the graft.



1. Select a section of stock with a flat surface, preferably on the south or southwest side so the prevailing wind will blow the graft into the tree instead of away. With a sharp knife, remove only the outer rough bark from an area large enough to conveniently accommodate the scion. Leave the bark sufficiently thick to hold the scion securely.



2. Choose a smooth, straight scion. Cut to about 6 inches long with 2 or 3 buds remaining. Make a smooth, straight sloping cut 2" to 3 1/2" long. Start this cut about 3" from the bottom of the scion on the opposite side and a little below a bud. This leaves an "insurance bud" in case the others fail to develop. Next, turn the scion over and make a chisel cut about 1/2" long on the back side at the lower end of the sloping cut. As an additional step, many grafters prefer to remove a thin layer of outside bark on both sides of the sloping cut, leaving a slight bevel. The bevel and the chisel cut provide additional cambium (uniting tissue) contact between the scion and the stock. The chisel cut also allows easier insertion of the scion. Keep hands free from the exposed areas.




3. & 4. Place prepared scion on smooth area of stock. Make pattern cuts through the bark down to the wood along both sides of the scion. The pattern cuts should be approximately half the length of the sloping cut. This produces the inlay pattern from which the name inlay bark graft is derived. Perform all the grafting operations as quickly, yet as accurately, as possible. When practical, work with your back to the sun to form a shade over the exposed areas.




5. Score the bark slightly, below and between the lower end of the pattern cuts. The score should not be completely through the bark as are the pattern cuts.




6. Loosen the flap (inlay) at the top with the tip of the knife blade.




7. Start chisel end of scion into the slot with the sloping cut against the wood of the stock.




8. Maintain pressure on the flap with the thumb while firmly forcing the scion into the slot. Do not allow the flap to loosen more than necessary to insert the scion.




9. Leave approximately 1/8" of the sloping cut above the top of the stock to encourage callus tissue development and to provide secure anchorage of the scion.




13. Use a medium-sized rubber band, masking tape, or grafting tape to secure the bag around the scion, at the cut corner just above the sloping cut.




10. Remove about half of the bark flap and nail the scion firmly to the stock as shown. If the scion is small, do not nail. One No. 18 wire nail 5/8" to 3/4" long, depending on the size of the scion, is usually sufficient. Some grafters use a staple applied with a hand staple gun instead of the nail. The staple must be long enough to go through the scion. For additional support, a second nail or staple may be placed below the first through the flap and scion and into the stock wood. Wrap securely with masking tape or grafting tape.




14. Tie the bag around the stock with string or masking tape to prevent its blowing off.



11. Take a piece of regular household aluminum foil, 8 by 12 inches, and wrap the upper 6" or 8" of the stock, covering the top also. Use the shiny side out. Permit the foil to extend up on the scion about 1/2". Press the foil firmly with the fingers until it remains in place.



12. Clip the corner of a quart-size polyethylene bag and carefully slip it down over the stock with the scion protruding through the clipped corner. Watch the buds to make sure they do not get damaged as bag is applied.



At this point, many grafters use amber shellac to coat the top and sides of the scion down to the polyethylene bag. Make sure the shellac is water based amber or orange shellac.

Another helpful strategy to protect the graft is attaching a bird perch below the graft that extends up above the graft union. The perch can be a bamboo stake or a stick that was removed from the rootstock. This provides birds with another area to sit without wiggling and loosening the graft union.

The aluminum foil, polyethylene bag, and amber shellac prevent excessive drying of the scion and exposed areas. The foil also reflects the sun's rays preventing excessive heat buildup around the graft union.

Aftercare of the Graft

The care of the tree during the first few years after grafting is as important as the operation itself.

A successful graft will usually begin growth in about three weeks, but can be slower if graft is performed early and the weather is cool. Grafts from two-year-old wood are usually slower to start growth than those from one-year-old wood.

Shoots from the graft may grow so rapidly that they become top-heavy and are easily damaged by wind. Such shoots should be supported by tying them to one-by-two inch stakes nailed to the stock. Excessive growth may be reduced by removing the growing tips of shoots after they have made a growth of 20 to 30 inches.

Some additional tips include:

- If the grafts are in an area being grazed, and within reach of livestock, install a protective fence or barrier around the tree.

- During the first summer, remove shoots that arise on the stock near the grafted area.
- Clear out other tree growth in the grove to give the grafted trees adequate growing space.
- If the graft is growing slowly, cut back some of the native branches left on the stock at the time of grafting. This will encourage additional growth of the graft.
- The poly bag and foil may be removed in late July or August.
- The square shoulder opposite the graft on large stocks (4 inches) will often dry and decay before healing. Healing can be accelerated by removing this edge of the stock with a sloping cut at a 20° to 30° angle downward and away from the side where the graft is growing. This operation can be done during the first winter.
- When two grafts are placed on one stock and both grow, the least desirable one should be removed during the first winter. The extra graft can be removed with a portion of

the stock as suggested above for removing the square shoulder on large diameter stocks. If both grafts are allowed to grow, a weak V-crotch angle will likely develop, causing the tree to be vulnerable to breaking and splitting.

About three years are required to grow a new top from the scion. During this time, the native branches should be removed. For example, if three native branches were left on the stock, remove one each year. This practice depends upon the vigor of the tree and graft. Delay removal of these native limbs if the grafts are making excessive growth. Likewise, hasten their removal if the graft is growing slowly. Remove the native branches flush with the main trunk.

Train the new tree to a central leader system by cutting back or removing side branches that crowd and compete with the leader. Consult HLA-6245 for more information on Training Pecan Trees.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0311 GH Revised.