



Johnsongrass in Pastures: Weed or Forage?

January 2017

Alex Rocateli
Forage System Extension Specialist

Misha Manuchehri
Weed Science Extension Specialist

Oklahoma Cooperative Extension Fact Sheets
are also available on our website at:
<http://osufacts.okstate.edu>

Johnsongrass (*Sorghum halepense* L.) is a warm-season grass. Originally from the Mediterranean region, it was introduced to North America in the 1800s as a forage alternative. The name “Johnsongrass” refers to Colonel William Johnson, who introduced this species to his river-bottom farm in Alabama in the 1840s. Today, Johnsongrass is found in all states except Minnesota, and is considered a noxious weed in 19 states (NRCS, 2016). Johnsongrass is popularly known as “the weed we love to hate and hate to love.” We “hate to love” it because it is one of the most common weeds in 30 different crops including corn, sorghum, cotton and soybeans. It also serves as a host for several insects, disease pathogens and nematodes of corn and sorghum. On the other hand, we “love to hate” it because it is not only a valuable forage due to its high yield, palatability and quality, but also is successful in reducing soil erosion as a plant cover alternative (Warwick and Black, 1983).

Identification

A Johnsongrass seedling can resemble a corn or sorghum seedling; however, its stems and leaves are narrower and completely hairless. In any developmental stage, Johnsongrass leaves have a very distinct and prominent white mid-vein, which differentiates it from most other grasses (Figure 2a). Adult plants can range in height from 2.5 to 7 feet tall. Johnsongrass is a bunch-type grass, tillering from



Figure 1. Established Johnsongrass along a fence line.




Figure 2. Visual clues to identify Johnsongrass.

the crown of the plant. The flower head is a panicle (highly branched) and can reach up to 1.5 feet long. It has a green central stalk (i.e., rachis) that contains several whorls of two to three lateral branches. The branches are perpendicular to the stem at the bottom and parallel at the top, resulting in a pyramidal-shaped head (Figure 2b). The seeds, which are very small (2 to 3 mm) and egg-shaped, attach to the lateral branches and turn from greenish-violet to dark reddish-brown when mature.

Both roots and rhizomes (i.e., horizontal underground stems) are found on Johnsongrass plants. Most other grass species that resemble Johnsongrass do not contain rhizomes. Rhizomes are white to brown in color and may contain purple spots and nodes covered by brown scaly sheaths (Figure 2c). The leaf collar, where the leaf sheath and leaf blade meet, can serve as a useful identification tool. Pulling the collar back and detaching it from the stem will reveal the presence of a toothed membrane called the ligule. With age, some ligules may develop a fringe of hairs in the upper portion (Figure 2d).

Life Cycle and Adaptability

Johnsongrass is an aggressive perennial. Either new shoots from rhizomes or new seedlings will sprout during early to mid-spring. Seeds start to germinate when soil temperatures reach 70 F; however, new shoots from rhizomes will sprout when soil temperatures are 60 F. Sprouts from rhizomes develop faster than seedlings by taking advantage of rhizome carbohydrates accumulated during the winter. Plants start to produce new rhizomes after five to seven true leaves have developed. This occurs approximately three to six weeks after emergence. Flowering will commence six to nine weeks after emergence, and viable seeds will be produced two to three weeks after flowering. During the fall, Johnsongrass growth ceases when soil temperatures return to 60 F, turning the plant dormant. In Oklahoma, Johnsongrass will start to grow by the end of March, and new rhizomes will start to develop by the end of April. Flowering will start in early June and viable seeds will appear in late June. Additionally, new rhizomes, flowers and seeds will continue to be produced until early November, when plants turn dormant.

Johnsongrass is adapted to a wide range of soil types within a pH range of 5 to 7.5. Therefore, Johnsongrass is mainly found in arable lands, orchards, open waste grounds, roadsides, pastures, irrigated canals and ditches. It grows best in fertile lowland soils. It is not adapted to poorly drained clay soils, but it can tolerate short periods of flooding. Rhizome production also is affected by soil type. Greater rhizome production and depth will occur in lighter-textured soils. For instance, clay soils will allow only half of the rhizomes that are capable of being produced in sandy loam soils. In addition, most rhizomes in clay and sandy loam soils will reach depths of 3 and 5 inches, respectively.

Johnsongrass as a Forage Alternative

Johnsongrass is very competitive and has desirable forage traits. It has relatively high quality and produces comparable yields (2 to 5 tons per acre) to other summer forages. Furthermore, Johnsongrass is highly palatable prior to reproductive growth. However, its palatability and quality quickly decrease after flowering, and cattle will avoid it. Table 1 compares forage dry yield, crude protein (CP), and total

Table 1. Forage dry yield, crude protein (CP) and total digestible nutrients (TDN) of summer forage crops.

<i>Summer forage crop</i>	<i>Dry yield (tons/A)</i>	<i>CP (%)</i>	<i>TDN (%)</i>
Bermudagrass, common	2-6	9-11	50-56
Bermudagrass, coastal	5-8	10-14	55-60
Johnsongrass	2-5	10-14	55-60
Sudangrass	2-6	9-12	55-60
Pearl millet	2-6	9-12	55-60
Foxtail millet	2-3	9-12	55-60
Sericea lespedeza	1-3	14-17	50-55

Source: Ball et. al (2007)

digestible nutrients (TDN) among Johnsongrass and other common forages. Values demonstrate that Johnsongrass, when correctly managed, can be a good forage option and comparable to bermudagrass and other common perennial forages (introduced and native).

Johnsongrass has some management considerations when used as a forage. The first is its high susceptibility to overgrazing. Cattle often kill Johnsongrass stands by grazing it to the ground. To maintain good Johnsongrass stands in pastures, start grazing or haying before flowering—when plants reach 12 to 18 inches—and stop when plants are grazed down to 6 to 8 inches. As previously discussed, rhizome energy will be very low before flowering; hence, grazing all of the leaves and stems at this stage will eventually starve the plant to death. Keeping a 6- to 8-inch stubble height also will maintain growing points and leaves, assuring regrowth back to 12 to 18 inches if a rest period of 30 to 45 days is allowed. This management strategy will not only help maintain forage quality, but also will maximize animal gain, avoid seed production and decrease infestations in neighboring field crops.

The second consideration is its high potential of nitrate and prussic acid accumulation. Both are poisonous—even lethal—to livestock at high levels. Nitrate and prussic acid poisoning occurs when plants resume growth after undergoing stress. For instance, Johnsongrass will accumulate nitrate or prussic acid after a rainfall that is followed by a severe drought period. The same challenges exist in other sorghums commonly used for hay or grazing. To avoid nitrate and prussic acid poisoning, use the following recommendations:

- test forage for nitrate and prussic acid concentrations after long drought periods or frost;
- wait at least five to seven days before grazing or cutting hay if high levels were detected;
- be extra cautious when grazing pastures recently fertilized with nitrogen; and
- *never introduce horses to Johnsongrass because it can result in equine cystitis.*

For more information on nitrate and prussic acid poisoning consult the factsheets PSS 2903 Nitrate Toxicity in Livestock (available at: <http://pods.dasnr.okstate.edu/docushare/dsweb/Get/Document-1996/PSS-2903web2013.pdf>) and PSS 2904 Prussic Acid Poisoning (available at: <http://pods.dasnr.okstate.edu/docushare/dsweb/Get/Document-6191/PSS-2904pod2013.pdf>).

Johnsongrass Management

Most producers prefer to control Johnsongrass in their pastures due to the management considerations previously described. Improved forages exist that have faster growth rates than Johnsongrass with no poisoning concerns. Prevention of Johnsongrass is the best line of defense against unwanted plants in pastures. Prevention practices include using weed-free seed, avoiding driving machinery through Johnsongrass stands, cleaning equipment after moving from an infested area and managing field margins. If Johnsongrass does become established; cultural, mechanical and chemical tools can be effective.

The easiest and most profitable way to control established Johnsongrass in pastures is to overgraze or continuously mow. Understanding the changes in rhizome reserves is key for management of Johnsongrass. Rhizome reserves will be minimal—from one to four weeks after emergence—as new sprouts are taking up all of the reserves for leaf and stem growth. Removing the leaves and stems at this stage will leave the plant without energy for regrowth. However, rhizome reserves will be reaching its maximum after flowering, when the plant starts to senesce. Removing leaves and stems at this stage will promote more leaf and stem growth if the weather is suitable for growth.

Mechanical control is another option, but is more expensive. Fall plowing to a depth just below the rhizomes (3 to 5

inches) is very effective because it will bring the rhizomes to the soil surface and expose them to killing temperatures. Disking has the opposite effect. It will increase infestations by cutting the rhizomes into pieces that will eventually sprout as new plants. However, combining disking with proper herbicide applications can effectively control Johnsongrass as wounds promote herbicide absorption. For the best results, make post-emergence applications when Johnsongrass is actively growing and is at least 18 to 24 inches tall and up to the heading stage. See Table 2 for herbicide control options. *Before applying any herbicide, always consult the label for appropriate rates, the addition of surfactants, tank-mix partners, plant-back intervals, geographic restrictions, spray carrier volume, nozzle selection and other special instructions.*

References

- Ball, D.M., C.S. Hoveland and G.D. Cacefield. 2007. *Southern Forages*, 4th ed. Potash and Phosphate Institute and the Foundation for Agronomic Research.
- Natural Resources Conservation Service (NRCS). 2016. PLANTS Database. Available at: <http://plants.usda.gov/core/profile?symbol=SOHA>. Data retrieved on 11/17/2016. United States Department of Agriculture.
- Warwick S.I. and L.D. Black. The Biology of Canadian Weeds. 61. *Sorghum halepense* (L.) PERS. *Can. J. Plant Sci.* 63: 997-1014.

Table 2. Herbicide options for Johnsongrass control in pasture.

<i>Herbicide Trade Name</i>	<i>Herbicide Common Name</i>	<i>Manufacturer</i>	<i>Mode of Action</i>	<i>Application Timing</i>	<i>Labeled Crops</i>	<i>Rate</i>	<i>Grazing Restrictions</i>
Balan DF™	Benefin	Trademark of Dow AgroSciences	Root Inhibitor	PPI ^a	Alfalfa and clover	2 to 2.5 lbs/A	No grazing restrictions
Outrider®	Sulfosulfuron	Monsanto	ALS Inhibitor	POST	Bermudagrass, Bahiagrass, and pastures west of the Mississippi River	0.75 to 2 oz/A	No grazing restrictions
Pastora®	Nicosulfuron Metsulfuron	Trademark of Dow AgroSciences and Bayer CropScience	ALS Inhibitor ALS Inhibitor	POST	Bermudagrass	1 to 1.5 oz/A	Do not graze until plants are dry
Plateau®	Imazapic	BASF	ALS Inhibitor	POST	All Pastures	2 to 12 oz/A	No grazing restrictions
Roundup PowerMAX®	Glyphosate	Monsanto Synthesis Inhibitor	EPSP Inhibitor	Preplant, PRE, PR, ST, WA and POST	Alfalfa (RR), bermudagrass, fescue, winter wheat, and others	8 to 44 fl oz/A	Restrictions dependent on application timing. READ LABEL

^a Abbreviations: PPI=preplant incorporated, POST=postemergence, PRE=preemergence, PR=pasture renovation, ST=spot treat, WA; wiper applicator.

The Oklahoma Cooperative Extension Service

WE ARE OKLAHOMA

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
 - It is administered by the land-grant university as designated by the state legislature through an Extension director.
 - Extension programs are nonpolitical, objective, and research-based information.
 - It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
 - More than a million volunteers help multiply the impact of the Extension professional staff.
 - It dispenses no funds to the public.
 - It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
 - Local programs are developed and carried out in full recognition of national problems and goals.
 - The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
 - Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies; Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 42 cents per copy, 0117 GH.