



Field Key to Larvae in Alfalfa

Phillip G. Mulder
Professor of Entomology

Richard Grantham
Director, Plant Disease and Insect Diagnostics Laboratory

Don C. Arnold
Retired Survey Entomologist

This key is designed to serve as a guide to identification of the more typical larvae of the common insect species found in the alfalfa fields of Oklahoma during the mid to late season. A 10 to 15 power hand lens will be most helpful in using this key. The identifying characters used are based upon those found on full-grown or nearly full-grown larvae and may not necessarily occur on newly hatched larvae. If the larva in question does not fit the proper identification furnished, recheck the specimen with the key. If it continues to key out improperly or is not one of the species listed, and proper identification is desired, place the larva in a small bottle containing 70 percent alcohol and mail to: Department of Entomology, Oklahoma State University, Stillwater, Oklahoma. Please do not send specimens for identification unless they are causing or suspected of causing damage to the crop. Please include information as to the type and amount of damage noted as well as the date and community where the larvae were collected. This information will assist in getting a more accurate and rapid reply to your questions.

Some insects found in alfalfa fields cannot be identified with this key. This would include adult insects, arthropods other than insects, and such insects as aphids and lygus bugs that do not have a larval stage. Be sure you have an insect larvae before attempting to use this key.

Occasional early season pests, such as cutworms, have not been included in the key as they are not normally serious in Oklahoma. If found, they should run to the last couplet, species not included in the key. If found causing serious damage, please send in specimens for identification.

This key should not be used for larvae occurring in crops other than alfalfa. Other keys are available for other crops and can be obtained from the local county extension center.

Survey Methods

Insect counts in alfalfa are taken on a per 10 sweep basis during the actively growing season. Using a standard 15-inch sweep net, a sweep consists of a half circle or a 180 degree arc allowing the net to sweep into the foliage as the surveyor walks across the field. A sweep is taken with every one or two steps. To get a good overall count, 10 sweeps should be taken in at least 5 well spaced places in the field. Numbers for each species should be averaged and reported as the number per

Oklahoma Cooperative Extension Fact Sheets
are also available on our website at:
<http://osufacts.okstate.edu>

10 sweeps. Larger populations (aphids, etc.) can be reported as the number per sweep.

Newly cut or overwintering alfalfa infestations are recorded as the number per square foot of crown. A wire form containing 1 square foot can be used as a measurement gauge. All insects within this area are counted and recorded as the number of insects per square foot. Depending on the size of the field, 3 to 5 counts should be taken in each field.

Newly planted alfalfa should be checked on a per linear foot basis. Count and report the number of insects present on the plants in a foot of drill row. Counts should be made in at least 5 locations throughout the field.


Descriptions of Larvae


Corn Earworm (*Heliothis zea*) The main distinguishing characteristic of this species is the distinct, short, sharp microspines, resembling whiskers, which are present between the longer hairs on the back. This gives the larva an "unshaven" appearance when viewed with a 10 to 15 power hand lens. (Do not confuse the pebbled or granular skin of other larvae with the microspines.) The body color varies greatly from light to dark green, pink, or brownish-yellow. When fully grown, the larvae measure up to 1½ inches in length. This destructive pest causes damage by feeding on the foliage.

Webworms (Several spp.) This may be one of three closely related species, the garden webworm (*Achyra rantalis*), the alfalfa webworm (*Loxostege commixtalis*), or the beet webworm (*Loxostege sticticalis*), which **can not** be easily separated in the field. Although the control is the same, please send in specimens for identification to insure accurate reporting. All of these larvae are greenish-yellow in color with three distinct, elongate, black spots arranged in a triangle on the upper portion of the side of each segment with the exception of the front three. The fully grown larvae measure about 1 inch in length. These larvae are foliage feeders and may move into alfalfa fields from other nearby crops or weeds. They usually spin webs over the leaves upon which they are feeding, hence the common name "webworms."

Green Cloverworm (*Plathypena scabra*) This is the only species commonly found in alfalfa that has four pairs of prolegs. The body color is green. The younger larvae have two longitudinal white stripes on each side, but these are absent

A Field Key to Some Common Larvae found in Alfalfa in Oklahoma





Prepared by
Horace W. Van Cleave
Revised by
Don C. Arnold
Entomology Department
Oklahoma State University

in the last instar. They may be up to 1 inch long when fully grown. They are primarily foliage feeders.

Alfalfa Caterpillar (*Colius eurytheme*) These larvae are covered with many short hairs. The body color is green with a narrow white stripe down each side. Inside each stripe is a very fine red line. They measure up to 1½ inches in length when fully grown. They are foliage feeders.

Cabbage Looper (*Trichoplusia ni*) These larvae move in a characteristic “looping” manner. They are larger at the back end and taper toward the head. The body is green with narrow white lines running the length of the body and is without black spots. These larvae are up to 1⅔ inches in length when fully grown. They are usually found feeding on leaves, giving the foliage a ragged appearance.

Fall Armyworm (*Spodoptera frugiperda*) These larvae usually have a distinct, broad, white inverted “Y” present on the front of the head (not to be confused with a narrow inverted “V” found on a few other species). The body varies from light tan to green to dark brown or nearly black in color with three widely separated narrow yellowish-white stripes down the back. On each side are three more broad longitudinal lines side by side with the top, brown; the middle, reddish; and the bottom, yellow with reddish mottlings. These larvae measure up to 1⅓ inches in length when fully grown. They are primarily foliage feeders.

Woolly Worms (Family Arctiidae) This may be one of several members of this family. The most common one in the state is the salt-marsh caterpillar (*Estigmene acaea*), which is covered with long black, brown, or yellowish hairs. The larvae of this species may become almost 2 inches in length when fully grown. The pests in this family are primarily foliage feeders. If found causing serious damage, send in specimens for identification.

Measuring Worms (Family Geometridae) These larvae may have two or three pairs of prolegs. Those with three pairs are usually brown or yellowish brown with dark stripes. They

may measure 1 inch or more when fully grown. They are foliage feeders.

Yellowstriped Armyworm (*Spodoptera ornithogalli*) These larvae vary in color from pale gray to jet black, but all will have two yellow stripes down the back. The gray individuals have two narrow dark triangles on the back of each segment, but these cannot be distinguished in the darker forms. The head is mostly brown. These larvae measure up to 1⅓ inches in length when fully grown. They are primarily foliage feeders.

Weevil Larvae (Family Curculionidae) These larvae are whitish or yellowish, C-shaped, more or less cylindrical, and legless.

Lady Beetle Larvae (Family Coccinellidae) The body color is generally dark with bright yellow, orange, or red markings. The body is covered with numerous spines. In a few species, the body is covered with a waxy secretion and resemble mealybugs, but a check of the mouthparts will clear up the confusion (mealybugs have piercing-sucking mouthparts while lady beetle larvae have biting mouthparts). This group is highly beneficial with both the larvae and adults feeding on aphids, spider mites, eggs, and young of many other pests.

Syrphid Fly Larvae (Family Syrphidae) This may be one of several members of this family. They are cylindrical and somewhat peg-shaped with protuberances on most segments. They usually are light green in color and the entire body is covered with short, colorless microspines. They are found crawling about on the plants feeding on aphids.

Aphid-Lions (Family Chrysopidae) These small, active, light brown larvae measure up to ½ inch in length when full grown. Both the larvae (aphid-lions) and adults (lacewing flies) are beneficial since they feed upon aphids or plant lice and other small pests. Be sure that the specimen suspected of being in this group have biting mouthparts. There are several other groups, such as true bugs, Order Hemiptera, which are similar in body shape, but differ from them by having piercing-sucking mouthparts.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, the Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0413 Revised. GH