



# Pasture Recovery Following Drought

Daren D. Redfearn

Extension Forage and Pasture Management Specialist

## Damage Assessment and Recovery

Following drought, stand damage is readily apparent even on well-managed pastures. When drought occurs during active growth of warm-season grass pastures, it is almost certain that root growth will be restricted, in addition to the more obvious decline in forage yield. Extent of stand damage due to differences in soil types, fertility practices, grazing management, pasture species, and harvest management is somewhat easy to determine.

### Soil type, fertility, grazing management

Pastures on lighter soils will generally show greater damage than stands on heavier soils. Pastures with lower fertility levels that were overgrazed will likely have greater damage than non-grazed pastures.

### Pasture species


Following drought, summer pastures will exhibit varying degrees of damage. Recently established stands with sparse growth may have few plants present. Established pastures that were grazed for the majority of the summer can have moderate to severe degrees of damage. This damage will be almost entirely due to stocking rate. On the other hand, pastures managed solely for hay production should be less damaged than grazed pastures. This is due to longer growth periods during the drought that excluded grazing for several months during the drought.

### Stand damage

Slightly damaged stands (less than 30 percent stand loss) should recover quickly with weed control, proper fertility, and deferred grazing or harvest once satisfactory growing conditions return. Stands that are moderately damaged (between 30 percent and 60 percent stand loss) should fully recover with weed control, proper fertility, and deferred grazing or harvest (**Figure 1**). There are many tillers and seed that will aid in stand recovery. If these stands remain thin into the fall, it may be preferable to overseed with an adapted pasture legume or winter annual grasses. Severely damaged stands (greater than 60 percent stand loss) are going to require an enormous level of patience for adequate recovery. If stands are thin in the fall, it may be preferable to overseed legumes or winter annual grasses.

All perennial grasses have growth mechanisms that contribute to recovery of previously established stands. Bermudagrass is unique in that it has stolons, in addition to rhizomes. Although most of the above-ground stolons are likely dead,

Oklahoma Cooperative Extension Fact Sheets are also available on our website at:  
<http://osufacts.okstate.edu>



**Figure 1. Old World bluestem pasture with moderate drought damage (between 30 percent and 60 percent stand loss). Slightly damaged stands (less than 30 percent stand loss) will have twice this many plants and severely damaged stands (greater than 60 percent stand loss) will have half this many plants.**

there are many rhizomes below ground in the plant crown that can produce new growth. The Old World bluestems and weeping lovegrass have rhizomes, but each of these plants also produces substantial seed in most years. Similar to the bermudagrass, the rhizomes of the Old World bluestems and weeping lovegrass will produce new growth. Much of the seed produced will be viable and capable of germinating and producing new plants if adequate moisture is present.

The best approach to post-drought pasture management is to reduce competition and focus on moisture conservation and use. First and most importantly, it is necessary that aggressive weed control measures are used to reduce competition for moisture and nutrients. This allows the desirable plants the best opportunity for successful re-establishment of rhizomes and stolons in bermudagrass pastures and rhizomes along with germination of the seedbank in Old World bluestem and weeping lovegrass pastures. Secondly, follow fertilization based on a proper soil test. Since our recovery approach is to manage many of these pastures as a “new” seeding, it is important that our initial fertilization focus is on adding phosphorus (P) to stimulate root growth. Lastly, it is important to

prepare for grazing and harvest deferment. Depending on the severity of the pasture damage, it may be necessary to restrict grazing on some of these pastures for longer than three months.

## Approach for Post-drought Pasture Management

Many pastures appear as if there is no hope for reasonable recovery when conditions return to normal. The restricted root growth is more of a concern during and immediately following drought since these are the structures responsible for extracting moisture and nutrients from the soil. It is critical to focus on management practices that enhance root regrowth in order to facilitate pasture recovery.

Once favorable growing conditions return, there will remain lingering effects related to growth and competition. Without a doubt, there will be increased weed competition and pressure from both grassy and broadleaf weeds. Pasture growth will be slower, especially in overgrazed pastures due not only to weed competition, but also reduced energy reserves that promote early, vigorous spring growth. Even with substantial rainfall, recovery of warm-season grass pastures is going to be difficult to predict. Any pasture growth response to a management practice is going to be almost entirely dependent on available moisture.

The key to successful pasture recovery following a drought is patience. The best approach to managing pastures that were overgrazed due the drought is to manage them as new plantings. This includes aggressive weed control, fertility based on a proper soil test, and grazing deferment. The obvious question is, ***if I can manage only one practice, which one do I choose?*** The answer to this question is weed control. However, it is important to recognize that additional fertility coupled with deferred harvest will result in more rapid recovery than a single management practice alone.

### Weed control as a tool for post-drought recovery

Even though weed control is most important, it should not be made complicated by the number of commercially available pasture herbicides and premixes. The key to successful weed control is timeliness. From a forage production standpoint, it is best to spray a single time at an earlier date with a lower rate of herbicide, rather than too late with an increased herbicide rate.

The best control method for broadleaf weeds is with timely post-emergent herbicide application of products such as 2,4-D, dicamba, or other premixed pasture herbicides that contain these products. For pastures with perennial, problematic, or noxious weeds, a premix with additional herbicide active ingredients and a broader level of weed control may be necessary. Herbicides with broad spectrum control should be applied early when weeds are small. This reduces weed growth and competition and more importantly, conserves moisture for forage grass growth.

### Fertilization as a tool for post drought-recovery

Plant fertility needs should be given special consideration in drought-damaged pastures. Since many of these pastures were grazed repeatedly, it is likely that root growth will be restricted to the top few inches of soil. It is important that these pastures have a strong root system to speed the recovery of drought-damaged pastures.

### Bermudagrass pastures

Since drought-damaged stands should be managed as a new stand, plant P needs should be the primary consideration. Standard P fertility recommendations are adequate for stands with slight to moderate damage, but stands with severe damage require additional P, even when soil test results indicate adequate available P levels (**Table 1**).

Since bermudagrass pastures respond quickly to supplemental fertility, the addition of 50 lbs nitrogen (N) per acre will increase the recovery once the stolons have reached 6 to 10 inches in length. Nitrogen should only be applied if soil moisture is present. Once bermudagrass pastures begin to show signs of recovery, additional N can be applied to increase production if soil moisture is present. During drought or immediately following a drought, it is important that only enough N be applied for a 30-day production period based on the presence of adequate soil moisture. For example, this would be no more than 75 lbs N per acre per month.

**Table 1. Recommended phosphorus (P) application for drought-damaged bermudagrass pastures based on soil test.**

Soil test P	Stand damage		
	Severe	Moderate	Slight
	----- P <sub>2</sub> O <sub>5</sub> needed -----		
Low	75 lbs/acre	75 lbs/acre	75 lbs/acre
Moderate	60 lbs/acre	60 lbs/acre	60 lbs/acre
High	30 lbs/acre	20 lbs/acre	20 lbs/acre
Adequate	20 lbs/acre	0	0

### Old World bluestem and weeping lovegrass pastures

Neither Old World bluestem nor weeping lovegrass respond as quickly to fertilization as bermudagrass. However, each still have requirements for supplemental fertility. Similar to the recommendations for severely drought-damaged stands in bermudagrass, it is important to provide additional P when soil test results indicate adequate available P levels (**Table 2**). Additional N fertilization during the “new” establishment period for Old World bluestem and weeping lovegrass is not recommended since a small amount of N is present in many of the fertilizers used for P fertilization. During drought or immediately following a drought, it is important that additional N be applied only if adequate soil moisture is available. For example, this would be from 60 to no more than 75 lbs N per acre for the growing season.

### Grazing and harvest deferment as a tool for post-drought recovery

Since we are managing these drought damaged pastures as recent plantings, it is important that they have adequate time for establishment. The single most important strategy for pasture recovery will be to avoid the temptation to begin grazing as soon as plants begin their spring growth.

Overgrazing coupled with severe removal of top growth develops plants with shallow root systems. These plants need an opportunity to replenish their energy reserves and establish new root growth. It may be necessary for pastures that were

**Table 2. Recommended phosphorus (P) application for drought-damaged Old World bluestem and weeping lovegrass pastures based on soil test.**

Soil Soil test P	Stand damage		
	Severe	Moderate	Slight
	----- P <sub>2</sub> O <sub>5</sub> needed -----		
Low	60 lbs/acre	60 lbs/acre	60 lbs/acre
Moderate	40 lbs/acre	40 lbs/acre	40 lbs/acre
High	30 lbs/acre	20 lbs/acre	20 lbs/acre
Adequate	20 lbs/acre	0	0

used heavily during the drought to not be grazed during the entire growing season. In some cases, it may be necessary to use temporary fencing to allow for adequate grazing deferment. Ideally, spring and summer grazing of heavily grazed pastures should not begin until the plants have at least four or five weeks of growth.

It is important to maintain a critical level of stubble height. Under normal growing conditions, forage stubble height should be no less than 3 inches for bermudagrass and 5 inches for both Old World bluestem and weeping lovegrass. This serves two purposes. The first purpose is that the remaining leaf area provides the energy for plant growth. During drought recovery, this will increase root growth in addition to plant growth. The second purpose is for soil coverage. As the stubble breaks down, it forms a layer of litter on the soil surface. This improves soil moisture retention and also reduces soil temperature.

### Pasture Recovery Prescriptions

The two best measures of pasture recovery following drought are stand thickness and forage height. These characteristics are often independent of one another, which means that a pasture may be thin and tall or it may be short and thick. While either of these two scenarios can be adequate during periods with favorable growth conditions, pasture recovery will occur more quickly when pastures are both thick and tall. To enhance the post-drought pasture recovery, it is important to consider the following pasture management practices:

1. Save the best (pastures) for last. Since these pastures will recover the quickest, it is important that they be allowed to have unrestricted growth when the probability of moisture is greatest.
2. Pay special attention to P fertility needs first to increase root growth of drought-damaged pastures.

3. Don't graze or harvest too early. Following a drought, plants that appear healthy may still have a short root system.

### Recommendations

#### ***Warm-season grass pastures managed and harvested solely for hay***

Cool-season grassy weeds that are present can be grazed in winter or while the summer grass is dormant, glyphosate can be applied to control any remaining winter annual weeds with additional broadleaf weed control in the spring if needed. Following a proper soil test, any nutrient deficiencies, especially P, should be made prior to summer. Nitrogen fertilization can be applied during the summer based on available soil moisture.

#### ***Warm-season grass pastures with broadleaf weeds***

While the summer grass is dormant, glyphosate can be applied to control any winter annual weeds with additional broadleaf weed control in the spring if needed. Following a proper soil test, any nutrient deficiencies, especially P, should be made prior to summer. Nitrogen fertilization can be applied during the summer based on available soil moisture.

#### ***Warm-season grass pastures with grassy weeds***

Cool-season grassy weeds that are present can be grazed in winter or while the summer grass is dormant, glyphosate can be applied to control any remaining winter annual weeds with additional broadleaf weed control in the spring if needed. Following a proper soil test, any nutrient deficiencies, especially P, should be made prior to summer. Nitrogen fertilization can be applied during the summer based on available soil moisture.

#### ***Warm-season grass pastures with legumes (may or may not include broadleaf and grassy weeds)***

Cool-season grassy weeds that are present should be grazed during January. Fertilization with P and K fertilization to promote legume growth should be made in February through early March. If necessary, broadleaf weed control can be applied following legume production cycle. Nitrogen fertilization can be applied during the summer based on available soil moisture.

### Recovery Assessment

Pasture recovery following drought is difficult to predict. With inadequate precipitation, pasture recovery may show minimal progress even with proper management. Good growing conditions coupled with proper management practices may result in complete recovery for slightly- to moderately-damaged stands within one year. However, severely damaged stands may take longer than one year for recovery even with adequate moisture.

## The Oklahoma Cooperative Extension Service

### *Bringing the University to You!*

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0112 GH